

## AN EXTENSION OF THE CAYLEY–HAMILTON THEOREM FOR NONLINEAR TIME–VARYING SYSTEMS

TADEUSZ KACZOREK

Faculty of Electrical Engineering, Białystok Technical University  
ul. Wiejska 45 D, 15–351 Białystok, Poland  
e-mail: kaczorek@isep.pw.edu.pl

The classical Cayley-Hamilton theorem is extended to nonlinear time-varying systems with square and rectangular system matrices. It is shown that in both cases system matrices satisfy many equations with coefficients being the coefficients of characteristic polynomials of suitable square matrices. The proposed theorems are illustrated with numerical examples.

**Keywords:** extension, Cayley-Hamilton theorem, nonlinear, time-varying system

### 1. Introduction

The classical Cayley-Hamilton theorem (Gantmacher, 1974; Kaczorek, 1988; Lancaster, 1969) says that every square matrix satisfies its own characteristic equation. Let  $\mathbf{A} \in \mathbb{C}^{n \times n}$  (the set of  $n \times n$  complex matrices) and  $p(s) = \det[I_n s - \mathbf{A}] = \sum_{i=0}^n a^i s_i$ , ( $a_n = 1$ ) be the characteristic polynomial of  $\mathbf{A}$ . Then  $p(\mathbf{A}) = \sum_{i=0}^n a_i \mathbf{A}^i = \mathbf{0}_n$  (the  $n \times n$  zero matrix). The Cayley Hamilton theorem was extended to rectangular matrices (Kaczorek, 1988; 1995c), block matrices (Kaczorek, 1995b; Victoria, 1982), pairs of commuting matrices (Chang and Chan, 1992; Lewis, 1982; 1986; Kaczorek, 1988), pairs of block matrices (Kaczorek, 1988; 1998) as well as standard and singular two-dimensional linear (2-D) systems (Kaczorek, 1992; 1995a; Smart and Barnett, 1989; Theodoru, 1989). The Cayley-Hamilton theorem and its generalizations were used in control systems, electrical circuits, systems with delays, singular systems, 2-D linear systems, etc., cf. (Busłowicz, 1981; 1982; Kaczorek, 1992; 1994; Lewis, 1982; Mertzios and Christodolous, 1986).

In (Kaczorek, 2005a), the Cayley-Hamilton theorem was extended to  $n$ -dimensional ( $n$ -D) real polynomial matrices. An extension of the Cayley-Hamilton theorem for discrete-time and continuous-time linear systems with delay was given in (Busłowicz and Kaczorek, 2004; Kaczorek, 2005b).

In this paper, the Cayley-Hamilton theorem will be extended to the case of nonlinear time-varying systems with square and rectangular system matrices. To the best of the author's knowledge, the extension of the Cayley-Hamilton theorem for nonlinear time-varying systems has not been considered yet.

### 2. Square System Matrices

Consider the nonlinear time-varying system

$$\dot{x}(t) = \mathbf{A}(x, t)x(t) + \mathbf{B}(x, t)u(t), \quad (1)$$

where  $x(t) \in \mathbb{R}^n$  is the state vector,  $u(t) \in \mathbb{R}^m$  is the input vector and  $\mathbf{A} = \mathbf{A}(x, t) \in \mathbb{R}^{n \times n}$ ,  $\mathbf{B} = \mathbf{B}(x, t) \in \mathbb{R}^{n \times m}$ . The well-known notion of the characteristic polynomial (equation) for linear systems can be extended for nonlinear systems of the form (1) as follows.

**Definition 1.** *The polynomial*

$$\begin{aligned} p(s) &= \det [\mathbf{I}_n s - \mathbf{A}(x, t)] \\ &= s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0 \end{aligned} \quad (2)$$

with the coefficients  $a_k = a_k(x, t)$ ,  $k = 0, 1, \dots, n - 1$  depending on  $x$  and  $t$  is called the characteristic polynomial of the system (1). The equation  $p(s) = 0$  is called the characteristic equation of the system (1).

**Theorem 1.** *The system matrix  $\mathbf{A}(x, t)$  satisfies the equation*

$$\sum_{i=0}^n a_i \mathbf{A}^{i+k}(x, t) = \mathbf{0}_n, \quad k = 0, 1, \dots \quad (a_n = 1). \quad (3)$$

*Proof.* It is easy to check that

$$\begin{aligned} &[\mathbf{I}_n s - \mathbf{A}(x, t)] [\mathbf{I}_n s^{-1} + \mathbf{A}(x, t) s^{-2} \\ &+ \mathbf{A}^2(x, t) s^{-3} + \dots] = \mathbf{I}_n. \end{aligned} \quad (4)$$

Hence

$$\begin{aligned} [\mathbf{I}s - \mathbf{A}(x, t)]^{-1} &= \mathbf{I}_n s^{-1} + \mathbf{A}(x, t) s^{-2} \\ &\quad + \mathbf{A}^2(x, t) s^{-3} + \dots \end{aligned} \quad (5)$$

The substitution of (2) and (5) into the well-known equality (Gantmacher, 1974; Kaczorek, 1988):

$$\begin{aligned} \text{Adj}[\mathbf{I}_n s - \mathbf{A}(x, t)] \\ &= [\mathbf{I}_n s - \mathbf{A}(x, t)]^{-1} \det[\mathbf{I}_n s - \mathbf{A}(x, t)] \end{aligned}$$

yields

$$\begin{aligned} \text{Adj}[\mathbf{I}_n s - \mathbf{A}(x, t)] \\ &= [\mathbf{I}_n s^{-1} + \mathbf{A}(x, t) s^{-2} + \mathbf{A}^2(x, t) s^{-3} + \dots] \\ &\quad \times (s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0). \end{aligned} \quad (6)$$

Note that the adjoint matrix  $\text{Adj}[\mathbf{I}_n s - \mathbf{A}(x, t)]$  is a polynomial matrix in  $s$  (a matrix with a nonnegative power of  $s$ ).

Comparing the coefficient matrices at the same power  $s^{-(k+1)}$  of (6), we obtain (3). ■

**Remark 1.** For  $k = 0$ , from (3) we have the extension of the classical Cayley-Hamilton theorem for the nonlinear system (1):

$$\begin{aligned} p(\mathbf{A}) &= \mathbf{A}^n(x, t) + a_{n-1} \mathbf{A}^{n-1}(x, t) \\ &\quad + \dots + a_1 \mathbf{A}(x, t) + a_0 \mathbf{I}_n = \mathbf{0}_n. \end{aligned} \quad (7)$$

**Example 1.** Consider the nonlinear system (1) with

$$\mathbf{A} = \mathbf{A}(x, t) = \begin{bmatrix} x_1 e^{-t} & -2x_2^2 \\ x_1 e^{-t} & x_2^2 e^t \end{bmatrix}, \quad (8)$$

where  $x = [x_1 \ x_2]^T$ .

The characteristic polynomial of (8) has the form

$$\begin{aligned} \det[\mathbf{I}_n s - \mathbf{A}] &= \det \begin{bmatrix} s - x_1 e^{-t} & 2x_2^2 \\ -x_1 e^{-t} & s - x_2^2 e^t \end{bmatrix} \\ &= s^2 - (x_1 e^{-t} + x_2^2 e^t) s \\ &\quad + x_1 x_2^2 (1 + 2e^t). \end{aligned} \quad (9)$$

In this case,

$$a_1(x, t) = -(x_1 e^{-t} + x_2^2 e^t), \quad a_0(x, t) = (1 + 2e^t),$$

and using (3) we obtain, for  $k = 0$ ,

$$\begin{aligned} \begin{bmatrix} x_1 e^{-t} & -2x_2^2 \\ x_1 e^{-t} & x_2^2 e^t \end{bmatrix}^2 - (x_1 e^{-t} + x_2^2 e^t) \begin{bmatrix} x_1 e^{-t} & -2x_2^2 \\ x_1 e^{-t} & x_2^2 e^t \end{bmatrix} \\ + x_1 x_2^2 (1 + 2e^{-t}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} &= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \end{aligned}$$

and, for  $k = 1$ ,

$$\begin{aligned} \begin{bmatrix} x_1 e^{-t} & -2x_2^2 \\ x_1 e^{-t} & x_2^2 e^t \end{bmatrix}^3 - (x_1 e^{-t} + x_2^2 e^t) \begin{bmatrix} x_1 e^{-t} & -2x_2^2 \\ x_1 e^{-t} & x_2^2 e^t \end{bmatrix}^2 \\ + x_1 x_2^2 (1 + 2e^{-t}) \begin{bmatrix} x_1 e^{-t} & -2x_2^2 \\ x_1 e^{-t} & x_2^2 e^t \end{bmatrix} &= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

Therefore, the matrix (8) satisfies Eqn. (3) for  $k = 0, 1, \dots$  ♦

### 3. Rectangular System Matrices

Let us consider a rectangular matrix  $\mathbf{A}(x, t)$  with the number of its columns  $m$  greater than its number of rows  $n$ , i.e.  $m > n$ ,

$$\begin{aligned} \mathbf{A}(x, t) &= [\mathbf{A}_1(x, t) \ \mathbf{A}_2(x, t)] \in \mathbb{R}^{n \times m}, \\ \mathbf{A}_1(x, t) &\in \mathbb{R}^{n \times n}, \quad \mathbf{A}_2(x, t) \in \mathbb{R}^{n \times (m-n)}. \end{aligned} \quad (10)$$

Let

$$\begin{aligned} p_1(s) &= \det[\mathbf{I}_n s - \mathbf{A}_1(x, t)] \\ &= s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0, \end{aligned} \quad (11)$$

where the coefficients  $a_k = a_k(x, t)$ ,  $k = 0, 1, \dots, n-1$  depend on  $x$  and  $t$ .

**Theorem 2.** Let the characteristic polynomial of  $\mathbf{A}_1(x, t)$  have the form (11). Then the matrix (10) satisfies the equation

$$\begin{aligned} \sum_{i=0}^n a_i [\mathbf{A}_1^{m+i-n}(x, t), \mathbf{A}_1^{m+i-n-1}(x, t) \mathbf{A}_2(x, t)] \\ = \mathbf{0}_{nm}, \quad (a_n = 1), \end{aligned} \quad (12)$$

where  $\mathbf{0}_{nm}$  is the  $n \times n$  zero matrix.

*Proof.* By induction it is easy to show that

$$\begin{aligned} \begin{bmatrix} \mathbf{A}_1(x, t) & \mathbf{A}_2(x, t) \\ 0 & 0 \end{bmatrix}^k \\ = \begin{bmatrix} \mathbf{A}_1^k(x, t) & \mathbf{A}_1^{k-1}(x, t) \mathbf{A}_2(x, t) \\ 0 & 0 \end{bmatrix} \end{aligned} \quad (13)$$

for  $k = 0, 1, \dots$ . Using (11), we obtain

$$\det \begin{bmatrix} \mathbf{I}_n s - \mathbf{A}_1(x, t) & \mathbf{A}_2(x, t) \\ 0 & \mathbf{I}_n s \end{bmatrix} = s^{m-n} \det [\mathbf{I}_n s - \mathbf{A}_1(x, t)] = \sum_{i=0}^n a_i s^{m+i-n}. \quad (14)$$

From the classical Cayley-Hamilton theorem for the matrix

$$\begin{bmatrix} \mathbf{A}_1(x, t) & \mathbf{A}_2(x, t) \\ 0 & 0 \end{bmatrix}$$

we have

$$\sum_{i=0}^n a_i \begin{bmatrix} \mathbf{A}_1(x, t) & \mathbf{A}_2(x, t) \\ 0 & 0 \end{bmatrix}^{m+i-n} = \mathbf{0}_m \quad (a_n = 1). \quad (15)$$

The substitution of (13) into (15) yields

$$\sum_{i=0}^n a_i \begin{bmatrix} \mathbf{A}_1^{m+i-n}(x, t) & \mathbf{A}_1^{m+i-n-1}(x, t) \mathbf{A}_2(x, t) \\ 0 & 0 \end{bmatrix} = \mathbf{0}_m, \quad (a_n = 1) \quad (16)$$

Taking into account only the first  $n$  rows of (16), we obtain (12). ■

**Remark 2.** The matrix  $\mathbf{A}_1(x, t)$  can be constructed from any  $n$  columns of the matrix  $\mathbf{A}(x, t)$  (Kaczorek, 1988).

**Theorem 3.** Let the characteristic polynomial of  $\mathbf{A}_1(x, t)$  have the form (11). Then

$$\sum_{i=0}^n a_i \begin{bmatrix} \mathbf{A}_1^i(x, t) & \mathbf{A}_1^i(x, t) \mathbf{A}_2(x, t) \end{bmatrix} = \mathbf{0}_{nm}, \quad (a_n = 1). \quad (17)$$

*Proof.* From the classical Cayley-Hamilton theorem for the matrix  $\mathbf{A}_1(x, t)$  we have

$$\sum_{i=0}^n a_i \mathbf{A}_1^i(x, t) = \mathbf{0}_n, \quad (a_n = 1). \quad (18)$$

The postmultiplication of (18) by the matrix  $[\mathbf{I}_n \ \mathbf{A}_2(x, t)]$  yields (17). ■

**Example 2.** Consider the rectangular matrix

$$\mathbf{A}(x, t) = [\mathbf{A}_1(x, t) \ \mathbf{A}_2(x, t)] = \begin{bmatrix} e^{-t} \sin x_1 & e^{-2t} \cos x_2 & x_2 \sin x_1 \\ -e^t \cos x_2 & \sin x_1 & x_1 e^{-t} \end{bmatrix}, \quad (19)$$

where  $x = [x_1 \ x_2]^T$ .

The characteristic polynomial of the matrix  $\mathbf{A}_1(x, t)$  has the form

$$\begin{aligned} p_1(s) &= \det [\mathbf{I}_n s - \mathbf{A}_1(x, t)] \\ &= \det \begin{bmatrix} s - e^{-t} \sin x_1 & -e^{-2t} \cos x_2 \\ e^t \cos x_2 & s - \sin x_1 \end{bmatrix} \\ &= s^2 + a_1 s + a_0, \end{aligned} \quad (20)$$

where

$$\begin{aligned} a_1 &= a_1(x, t) = -(1 + e^{-t}) \sin x_1, \\ a_0 &= a_0(x, t) = e^{-t} (\sin^2 x_1 + \cos^2 x_2). \end{aligned}$$

Using (12), we obtain the result given by (21). Equation (17) in this case has the form (22). Therefore, the matrix (19) satisfies (12) and (17). ♦

If  $n > m$ , then the matrix  $\mathbf{A}(x, t)$  can be written in the form

$$\mathbf{A}(x, t) = \begin{bmatrix} \mathbf{A}_1(x, t) \\ \mathbf{A}_2(x, t) \end{bmatrix} \in \mathbb{R}^{n \times m}, \quad (23)$$

$$\mathbf{A}_1(x, t) \in \mathbb{R}^{m \times m}, \quad \mathbf{A}_2(x, t) \in \mathbb{R}^{(n-m) \times m}.$$

In much the same way as Theorem 1, the following dual theorem can be proved.

**Theorem 4.** Let  $\mathbf{A}(x, t)$  have the form (23) and

$$\begin{aligned} \bar{p}_1(s) &= \det [\mathbf{I}_n s - \mathbf{A}_1(x, t)] \\ &= s^m + \bar{a}_{m-1} s^{m-1} + \dots + \bar{a}_1 s + \bar{a}_0, \end{aligned} \quad (24)$$

where the coefficients  $\bar{a}_i = \bar{a}_i(x, t)$ ,  $i = 0, 1, \dots, m - 1$  are functions of  $x$  and  $t$ . Then

$$\sum_{i=0}^m \bar{a}_i \begin{bmatrix} \mathbf{A}_1^{n+i-m}(x, t) \\ \mathbf{A}_2(x, t) \mathbf{A}_1^{n+i-m+1}(x, t) \end{bmatrix} = \mathbf{0}_{nm} \quad (a_m = 1). \quad (25)$$

From the classical Cayley-Hamilton theorem for  $\mathbf{A}_1(x, t)$  and (24), we have

$$\sum_{i=0}^m \bar{a}_i \mathbf{A}_1^i(x, t) = \mathbf{0} \quad (\bar{a}_m = 1). \quad (26)$$

The premultiplication of (26) by the matrix

$$\begin{bmatrix} \mathbf{I}_m \\ \mathbf{A}_2(x, t) \end{bmatrix}$$

$$\begin{aligned}
 & \left[ \mathbf{A}_1^3(x, t), \mathbf{A}_1^2(x, t)\mathbf{A}_2(x, t) \right] + a_1(x, t) \left[ \mathbf{A}_1^2(x, t), \mathbf{A}_1(x, t)\mathbf{A}_2(x, t) \right] + a_0(x, t) \left[ \mathbf{A}_1(x, t), \mathbf{A}_2(x, t) \right] \\
 &= \left[ \begin{array}{l} e^{-3t} \sin^3 x_1 - 2e^{-2t} \sin x_1 \cos^2 x_2 - e^{-t} \sin x_1 \cos^2 x_2 \\ \cos^3 x_2 - e^{-t} \sin^2 x_1 \cos x_2 - e^{-2t} \sin^2 x_1 \cos x_2 - e^t \sin^2 x_1 \cos x_2 \\ \\ e^{-4t} \sin^2 x_1 \cos x_2 - e^{-3t} (\cos^3 x_2 - \sin^2 x_1 \cos x_2) + e^{-2t} \sin^2 x_1 \cos x_2 \\ - (e^{-2t} + e^{-3t}) \sin x_1 \cos^2 x_2 - e^{-t} \sin x_1 \cos^2 x_2 + \sin^3 x_1 \\ \\ e^{-4t} x_1 \sin x_1 \cos x_2 + e^{-3t} x_1 \sin x_1 \cos x_2 + e^{-2t} x_2 \sin^3 x_1 - e^{-t} x_2 \sin x_1 \cos^2 x_2 \\ e^{-t} x_1 \sin^2 x_1 - e^{-2t} x_1 \cos^2 x_2 - e^t x_2 \sin^2 x_1 \cos x_2 - x_2 \sin^2 x_1 \cos x_2 \end{array} \right] \quad (21) \\
 & - (1 + e^{-t}) \sin x_1 \left[ \begin{array}{ll} e^{-2t} \sin^2 x_1 - e^{-t} \cos^2 x_2 & e^{-3t} \sin x_1 \cos x_2 + e^{-2t} \sin x_1 \cos x_2 \\ - \sin x_1 \cos x_2 - e^{-t} \sin x_1 \cos x_2 & \sin^2 x_1 - e^{-t} \cos^2 x_2 \\ e^{-3t} x_1 \cos x_2 + e^{-t} x_2 \sin^2 x_1 & \\ e^{-t} x_1 \sin x_1 - e^{-t} x_2 \sin x_1 \cos x_2 & \end{array} \right] \\
 & + e^{-t} (\sin^2 x_1 + \cos^2 x_2) \left[ \begin{array}{lll} e^{-t} \sin x_1 & e^{-2t} \cos x_2 & x_2 \sin x_1 \\ -e^t \cos x_2 & \sin x_1 & x_1 e^{-t} \end{array} \right] = \left[ \begin{array}{lll} 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 & \left[ \mathbf{A}_1^2(x, t), \mathbf{A}_1^2(x, t)\mathbf{A}_2(x, t) \right] + a_1(x, t) \left[ \mathbf{A}_1(x, t), \mathbf{A}_1(x, t)\mathbf{A}_2(x, t) \right] + a_0(x, t) \left[ \mathbf{I}_n, \mathbf{A}_2(x, t) \right] \\
 &= \left[ \begin{array}{ll} e^{-2t} \sin^2 x_1 - e^{-t} \cos^2 x_2 & e^{-3t} \sin x_1 \cos x_2 + e^{-2t} \sin x_1 \cos x_2 \\ - \sin x_1 \cos x_2 - e^t \sin x_1 \cos x_2 & e^{-t} \cos^2 x_2 + \sin^2 x_1 \end{array} \right] \\
 & \left[ \begin{array}{l} e^{-4t} x_1 \sin x_1 \cos x_2 + e^{-3t} \sin x_1 \cos x_2 + e^{-2t} x_2 \sin^3 x_1 - e^{-t} x_2 \sin x_1 \cos^2 x_2 \\ e^{-t} x_1 \sin^2 x_1 - e^{-2t} x_1 \cos^2 x_2 - e^{-t} x_2 \sin^2 x_1 \cos x_2 - x_2 \sin^2 x_1 \cos x_2 \end{array} \right] \quad (22) \\
 & - (1 + e^{-t}) \sin x_1 \left[ \begin{array}{lll} e^{-t} \sin x_1 & e^{-2t} \cos x_2 & e^{-t} x_2 \sin^2 x_1 + e^{-3t} x_1 \cos x_2 \\ -e^t \cos x_2 & \sin x_1 & e^{-t} x_1 \sin x_1 - e^t x_2 \sin x_1 \cos x_2 \end{array} \right] \\
 & + e^{-t} (\sin^2 x_1 + \cos^2 x_2) \left[ \begin{array}{lll} 1 & 0 & x_2 \sin x_1 \\ 0 & 1 & x_1 e^{-t} \end{array} \right] = \left[ \begin{array}{lll} 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right].
 \end{aligned}$$

yields

$$\sum_{i=0}^m \bar{a}_i \left[ \begin{array}{l} \mathbf{A}_1^i(x, t) \\ \mathbf{A}_2(x, t)\mathbf{A}_1^i(x, t) \end{array} \right] = \mathbf{0}_{mn} \quad (\bar{a}_m = 1). \quad (27)$$

Therefore we have proved the following theorem.

**Theorem 5.** *Let the characteristic polynomial of  $\mathbf{A}_1(x, t)$  have the form (24). Then the matrix (23) satisfies Eqn. (27).*

**Remark 3.** Equation (12) can be obtained by the postmultiplication of (18) by the matrix

$$\left[ \mathbf{A}_1^{m-n}(x, t) \quad \mathbf{A}_1^{m-n-1}(x, t)\mathbf{A}_2(x, t) \right]$$

and Eqn. (25) by the premultiplication of (26) by the matrix

$$\left[ \begin{array}{l} \mathbf{A}_1^{n-m}(x, t) \\ \mathbf{A}_2(x, t)\mathbf{A}_1^{n-m-1}(x, t) \end{array} \right].$$

#### 4. Concluding Remarks

The Cayley-Hamilton theorem has been extended for nonlinear time-varying systems with square (Theorem 1) and rectangular (Theorems 2–5) system matrices. It was shown that in both cases the system matrices satisfy many equations. For rectangular system matrices, starting from characteristic polynomials of square matrices, it is possible to obtain many different equations that are satisfied by these system matrices. Note that the equations are satisfied for all parameters of nonlinear systems.

The presented generalizations can be extended to block matrices and two-dimensional nonlinear time-varying systems.

#### Acknowledgments

I wish to thank very much Professors K. Gałkowski and M. Busłowicz for their valuable suggestions and remarks. This work has been supported by the Polish Ministry of Education and Science, Grant 3T10A 066 27.

#### References

- Busłowicz M. (1981): *An algorithm of determination of the quasi-polynomial of multivariable time-invariant linear system with delays based on state equations*. — *Archive of Automatics and Telemechanics*, Vol. XXXVI, No. 1, pp. 125–131, (in Polish).
- Busłowicz M. (1982): *Inversion of characteristic matrix of the time-delay systems of neural type*. — *Found. Contr. Eng.*, Vol. 7, No. 4, pp. 195–210.
- Busłowicz M. and Kaczorek T. (2004): *Rechability and minimum energy control of positive linear discrete-time systems with one delay*. — *Proc. 12th Mediterranean Conf. Control and Automation*, Kasadasi-Izmir, Turkey (on CD-ROM).
- Chang F.R. and Chan C.N. (1992): *The generalized Cayley-Hamilton theorem for standard pencils*. — *Syst. Contr. Lett.*, Vol. 18, No. 3, pp. 179–182.
- Gantmacher F.R. (1974): *The Theory of Matrices*. — Vol. 2, Chelsea.
- Kaczorek T. (1988): *Vectors and Matrices in Automation and Electrotechnics*. — Warsaw: Polish Scientific Publishers, (in Polish).
- Kaczorek T. (1992–1993): *Linear Control Systems, Vols. I and II*. — Taunton: Research Studies Press.
- Kaczorek T. (1994): *Extensions of the Cayley-Hamilton theorem for 2D continuous-discrete linear systems*. — *Appl. Math. Comput. Sci.*, Vol. 4, No. 4, pp. 507–515.
- Kaczorek T. (1995a): *An existence of the Cayley-Hamilton theorem for singular 2D linear systems with non-square matrices*. — *Bull. Pol. Acad. Techn. Sci.*, Vol. 43, No. 1, pp. 39–48.
- Kaczorek T. (1995b): *An existence of the Cayley-Hamilton theorem for nonsquare block matrices*. — *Bull. Pol. Acad. Techn. Sci.*, Vol. 43, No. 1, pp. 49–56.
- Kaczorek T. (1995c): *Generalization of the Cayley-Hamilton theorem for nonsquare matrices*. — *Proc. Int. Conf. Fundamentals of Electrotechnics and Circuit Theory XVIII-SPETO*, Gliwice, Poland, pp. 77–83.
- Kaczorek T. (1998): *An extension of the Cayley-Hamilton theorem for a standard pair of block matrices*. — *Appl. Math. Com. Sci.*, Vol. 8, No. 3, pp. 511–516.
- Kaczorek T. (2005a): *Generalization of Cayley-Hamilton theorem for n-D polynomial matrices*. — *IEEE Trans. Automat. Contr.*, Vol. 50, No. 5, pp. 671–674.
- Kaczorek T. (2005b): *Extension of the Cayley-Hamilton theorem for continuous-time systems with delays*. — *Appl. Math. Comp. Sci.*, Vol. 15, No. 2, pp. 231–234.
- Lancaster P. (1969): *Theory of Matrices*. — New York: Acad. Press.
- Lewis F.L. (1982): *Cayley-Hamilton theorem and Fadeev's method for the matrix pencil  $[sE - A]$* . — *Proc. 22nd IEEE Conf. Decision and Control*, San Diego, USA, pp. 1282–1288.
- Lewis F.L. (1986): *Further remarks on the Cayley-Hamilton theorem and Fadeev's method for the matrix pencil  $[sE - A]$* . — *IEEE Trans. Automat. Contr.*, Vol. 31, No. 7, pp. 869–870.
- Mertizios B.G. and Christodolous M.A. (1986): *On the generalized Cayley-Hamilton theorem*. — *IEEE Trans. Automat. Contr.*, Vol. 31, No. 1, pp. 156–157.
- Smart N.M. and Barnett S. (1989): *The algebra of matrices in n-dimensional systems*. — *IMA J. Math. Contr. Inf.*, Vol. 6, pp. 121–133.
- Theodoru N.J. (1989): *M-dimensional Cayley-Hamilton theorem*. — *IEEE Trans. Automat. Control*, Vol. AC-34, No. 5, pp. 563–565.
- Victoria J. (1982): *A block Cayley-Hamilton theorem*. — *Bull. Math. Soc. Sci. Math. Roum*, Vol. 26, No. 1, pp. 93–97.

Received: 10 August 2005

Revised: 15 September 2005